





SECTION-A

1. (d) Vitamin B₁₂ (1)
2. (c) About three times (1)
3. (b) (1)
- 
4. (d) 40 min (1)
5. (b) Sorbitol (1)
6. (a) CC1=CC=CC=C1C#N (1)
7. (d) [Cr (H₂O)₆]Cl₃ (1)
8. (d) Benzyl alcohol (1)
9. (b) CrO₄²⁻ (1)
- 10.(b) Diethyl ether (1)
- 11.(a) CH₃ NH₂ (1)
- 12.(d) Aspirin (1)
- 13.(d) P-Benzoquinone (1)
- 14.(c) i_x = i_y = i_z (1)
- 15.(c) Assertion (A) True, Reason (R) False (1)
- 16.(b) Assertion (A) True, Reason(R) True (1)
- But Reason(R) not true explanation
- 17.(c) Assertion (A) True, Reason (R) False (1)
18. (d) Assertion (A) False, Reason (R) True (1)

SECTION-B

- 19.(a) Aniline being lewis base react with Anhydrous AlCl₃ which is lewis acid to form salt. (1)

(b) Methylamine accept proton from water and liberate OH^- ion which combine with Fe^{3+} ion to form hydrated ferric oxide $\text{Fe}(\text{OH})_3$ or (1)



OR

(a) Because Grignard reagent reacts with moisture and form Alkane. (1)

(b) C-Cl bond in chloro benzene acquire some double bond character due to delocalization of ions pair on chlorine so bond length decreases


OR

any other relevant answer. (1)


21. (a) amylose is water soluble linear polymer of α -D glucose whereas amylopectin is water insoluble branched (C_1 - C_6) glycosidic linkage carrying branched polymer. (1)

(b) Intra molecular H-Bonding (1)

22. Geometrical Isomers (1)



Optical isomers (1)



23. $Q = I \times t$

$$= 0.5 \times 4 \times 60 \times 60$$

$$= 20 \times 360$$

$$= 7200 \text{ C} \quad (1)$$

96500 corresponds to $6.02 \times 10^{23} \text{ e}^-$

$$7200 \text{ C} \quad \text{gives} = \frac{6.02 \times 10^{23}}{96500} \times 7200$$

$$= 4.49 \times 10^{22} \text{ e}^- \quad (1)$$

24.(a)(i) Azeotropic mixture is type of liquid mixture having definite

composition and boiling like a pure liquid (1/2)

eg. 95.37% $\text{C}_2\text{H}_5\text{OH}$ + 4.63% H_2O (1/2)

OR

Any other relevant example

(ii) Solutions which have the same osmotic pressure at same temperature (1/2)

eg. 0.9% solution of pure NaCl is isotonic with RBC (1/2)

OR

Any other relevant example

OR

(b) If we have two completely miscible volatile liquid A and B having mole fraction x_A and x_B . Then at certain temperature partial pressures P_A and P_B and vapour pressure in pure state P_A° and P_B° are expressed as

$$P_A = P_A^\circ \cdot x_A$$

$$P_B = P_B^\circ \cdot x_B$$


$$P_T = P_A + P_B \quad (1/2)$$

$$P_T = P_A^\circ \cdot x_A + P_B^\circ \cdot x_B$$

$$P_T = P_A^\circ (1 - x_B) + P_B^\circ x_B$$

$$\text{when } x_A = 1 \quad P_T = P_A^\circ \cdot x_A$$

$$\text{when } x_B = 1 \quad P_T = P_B^\circ x_B \quad (1/2)$$





- 25.(i) E_a decrease (1)
(ii) No effect on ΔG (1)

SECTION-C

26. (a) It is the amide linkage present between – COOH group of one α amino acid and NH_2 group of other amino acid. (1)
- (b) When protein in native form is subjected to physical changes like change in temperature or pH then hydrogen bonds are broken, it loses its biological activity and all structures are destroyed and only primary structure remain intact. (1)
- (c) It is the sequence in which various α -amino acids present in a protein are linked to one another. (1)


OR

Amino acids contain acidic and basic group within same molecule. In aqueous solution they neutralize each other, carboxyl group loses a proton and amino group accept it. (1)



NH_3^+ group act as acid	Amphoteric react with acid and base	COO^- group act as base	(1)
--------------------------------------	---	-------------------------------------	-----

27.(a) $\text{CH}_3\text{CH}_2\text{COOH}$ (1)



28.(a) 1st order (1)

(b) min^{-1} (1)

(c) $t_{1/2} = \frac{0.693}{K}$ (1)

29. For AB_2

$$M_{\text{AB}_2} = \frac{Kf \cdot W_B \cdot 1000}{W_A \cdot \Delta T f} \quad (1/2)$$

$$= \frac{5.1 \times 1 \times 1000}{20 \times 2.3}$$

$$= 110.87\text{u} \quad (1/2)$$

$$M_{\text{AB}_4} = \frac{5.1 \times 1 \times 1000}{20 \times 1.3}$$

$$= 196.5\text{u}$$

Atomic mass of A = a and Atomic mass of B is b

$$\therefore a + 2b = 110.87 \quad (\text{i}) \quad (1)$$

$$a + 4b = 196.5 \quad (\text{ii})$$

(ii) - (i)

$$196.5 - 110.87 = a + 4b - a - 2b$$

$$85.63 = 2b \quad (1)$$

$$b = 42.815\text{u}$$

$$a+2b=110.87$$



$$a+2 \times 42.64=110.87$$

$$\therefore a = 110.87 - 85.28$$




$$= 25.59$$

i.e. atomic mass of A = 25.59 u

atomic mass of B = 42.64 u




OR




31.(a) The difference of energy between the two sets of a orbitals is called as crystal field splitting energy. (1)

(b)




(2)


OR



H_2O being weak ligand, don't cause pairing 5 unpaired e (1)



CN^- strong ligand, cause pairing so there is 1 unpaired e (1)



(c) $\Delta_o > P$ pairing occurs (1)

$\Delta_o < P$ No pairing occurs


32. (a) $\text{C}_6\text{H}_5\text{CHClC}_6\text{H}_5$ (1)

(b) 1-Bromo pentane > 2-Bromopentane > 2-Bromo-2-methylbutane (1)

(c) Allylic carbocation is stable (1)



(d) I^- is better leaving group than Cl^- (1)

OR



(d)  Allylic substitution (1)

33.(a) (i) $4 \text{FeOCr}_2\text{O}_3 + 8 \text{Na}_2\text{CO}_3 + 7\text{O}_2 \rightarrow 8\text{Na}_2\text{CrO}_4 + 2\text{Fe}_2\text{O}_3 + 8\text{CO}_2$ (1)

(ii) $2\text{Na}_2\text{CrO}_4 + \text{H}_2\text{SO}_4 \rightarrow \text{Na}_2\text{Cr}_2\text{O}_7 + \text{Na}_2\text{SO}_4 + \text{H}_2\text{O}$ (1)
(Conc)



OR





OR

any other relevant answer.

(b) Lanthanoids

Actinoids

(i) Electronic Configuration



(1)

(ii) Regular decrease in

Regular decrease in

(1)

size from left to

size from left to

right known as

right is known as

lanthanoid contraction

Actinoid contraction

(iii) \Rightarrow Lanthanoids react with dilute acid to liberate

\Rightarrow Actinoids are highly reactive in divided state

H_2 gas

\Rightarrow Form oxide and hydroxides of type $\text{M}_2\text{O}_3 / \text{M}(\text{OH})_3$

\Rightarrow React with boiling water to give mixture of oxide and hydride

\Rightarrow With C form carbides

\Rightarrow Attacked by HCl but the effect of HNO_3 is very small.

\Rightarrow With halogen form halides

\Rightarrow No action of alkalies

OR

any other relevant difference


34.(a) $E^\circ_{\text{cell}} = E^\circ_{\text{cathode}} - E^\circ_{\text{Anode}}$

$$= E^\circ_{\text{Cu}^{2+}/\text{Cu}} - E^\circ_{\text{Mg}^{2+}/\text{Mg}}$$

$$= 0.34 - (-2.36)$$

$$= 2.70\text{V}$$

(1/2)



$$E_{\text{cell}} = E^\circ_{\text{cell}} - \frac{0.0591}{2} \log \frac{[\text{Mg}^{2+}]}{[\text{Cu}^{2+}]}$$

$$= 2.70 - \frac{0.0591}{2} \log \frac{0.001}{0.0001}$$

$$\begin{aligned}
 &= 2.70 - 0.0295 \log 10 \\
 &= 2.70 - 0.0295 \times 1 \\
 &= 2.6705\text{V}
 \end{aligned}
 \tag{1}$$

(b) Because the number of ions per unit volume decreases. (2)


OR

(a) (i) During recharging, cell is operated like electrolytic cell.

(ii) Electrical energy is supplied to it from external source.

(iii) Electrode reactions are reverse of that of discharging. (1)


(iv) At cathode (Reduction) (1)



At Anode (oxidation)




Overall reaction



(b) $E^\circ \text{Zn}^{2+}/\text{Zn} = -0.76\text{V}$

$E^\circ \text{Cu}^{2+}/\text{Cu} = 0.34\text{V}$






$$E^\circ_{\text{cell}} = 0.34 - (-0.76)$$

$$= 1.10\text{V}$$

(c) E°_{cell} +ve means reaction is spontaneous and in this reaction zinc is oxidised \therefore we can't store CuSO_4 in zinc pot. (2)

35. (a) (1)






(a) Phenol gives violet colouration with neutral FeCl_3 solution but benzoic acid does not. (1)

OR

any other relevant test

(b) Acetaldehyde is more reactive towards nucleophilic addition reaction because of steric hindrance in acetone. (1)

(c) (1)



(1)