## **Model Question Paper**

Class-XII (Session: 2020-21) Subject-Mathematics ( Regular 2) Maximum Marks: 85 Time Allowed: 3 hrs **Special Instructions:-**Same as that of Previous Years Annual question paper March 2020. (i) Q1 to 10 are multiple choice questions and are of 1 mark each. Q. 11 to 13 are of 3 marks each. Q14 to 22 are of 4 marks each and Q. 23 to 27 are of 6 marks each. (ii) All questions are compulsory. (iii) 30% more internal choices have been provided from 70% of the syllabus, as 30% syllabus has been deleted due to COVID-19 Pandemic for the session 2020-21 only. 1.  $\cos^{-1}\left(\cos\frac{7\pi}{6}\right)$  equals to 1 (a)  $\frac{7\pi}{6}$  (b)  $\frac{5\pi}{6}$  (c)  $\frac{\pi}{3}$  (d)  $\frac{\pi}{6}$ 2. Let A be a nonsingular square matrix of order 3 × 3. Then |adjA| is equal to (c)  $|A|^3$ (d) 3|A| (b)  $|A|^2$ (a) |A| 3.  $\frac{d}{dx}(e^{-x})$  in equal to 1 (b)  $-e^{-x}$  (c)  $\frac{1}{e^x}$  (d)  $\frac{-1}{e^x}$ (a)  $e^{-x}$ 1 4. The function f(x) = Sinx is

- (a) Increasing in  $[0, \pi/2]$
- (b) Decreasing in  $[0, \pi/2]$
- (c) Neither increasing nor decreasing in  $[0, \pi/2]$
- (d) None of these

5. 
$$\frac{-1}{\sqrt{1-x^2}} dx$$

- (a)  $\sin^{-1} x + c$  (b)  $\cos^{-1} x + c$  (c)  $\tan^{-1} x$  (d)  $\tan^{-1} x + c$
- 6. The degree of the differential equations.  $(z'')^3 + (z')^2 + \operatorname{Sin}(z') + 1 = 0$
- (a) 3 (b) 2 (c) not defined (d) None of these
- 7. The direction ratios of the vector  $\hat{\mathbf{i}} + 2\hat{\mathbf{j}} + 3\hat{\mathbf{k}}$  is
  - (a) <1, 2, 3> (b) <2, 1, 3>
  - (c)  $<\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}>$  (d) <1, -2, 3>
- 8. If  $\theta$  be the angle between any two vectors  $\vec{a}$  and  $\vec{b}$  then  $|\vec{a} \vec{b}| = |\vec{a} \times \vec{b}|$  when  $\theta$  is equal to
  - (a) 0 (b)  $\frac{\pi}{4}$  (c)  $\frac{\pi}{2}$  (d)  $\frac{1}{\pi}$
- 9. If a line makes angles 90°, 135°, 45° with x, y and z=axis respectively then its direction cosines are
  - (a)  $<0, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}>$  (b)  $<0, \frac{-1}{\sqrt{2}}, 1>$
  - (c)  $<0, -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}>$  (d) none of these

10. If A and B are events such that P(A|B) = P(B|A) then

(a) A ⊂ B But A≠B

(b) A = B

(c)  $A \cap B = \phi$ 

(d) P(A) = P(B)

11. Find all points of discontinuity of function given by

 $f(x) = \begin{cases} x + 1 & \text{if } x \ge 1 \\ x^2 + 1 & \text{if } x < 1 \end{cases}$ 

Find  $\frac{dy}{dx}$  if  $y = \cos^{-1} \left[ \frac{1 - x^2}{1 + x^2} \right]$ , 0 < x < 1

12. Show that  $z = \log(1+x) - \frac{2x}{2+x}$ , x > -1 is an increasing functions of

x throughout its domain.

3

4

1

3

Or

Find the equation normal at the point (am<sup>2</sup>, am<sup>3</sup>) for the curve  $ay^2 = x^3$ 

13. Determine P (E|F) when Mother, Father and Son line up at random for the family picture.

E: Son on one end

F: Father in middle.

14. Let L be the set of all lines in a plane and R be the relation in L defined as  $R = \{L_1, L_2\}$ : L<sub>1</sub> is perpendicular to L<sub>2</sub>\}. Show that R is symmetric but neither reflexive nor transitive.

Or

Find go f and fog. If

$$f(x) = |x|, g(x) = |5x-2|$$

15. Prove that  $\tan^{-1} \sqrt{x} = \frac{1}{2} \cos^{-1} \left( \frac{1-x}{1+x} \right), x \in [0,1]$ 

Write in the simplest form.

$$\tan^{-1}\left(\frac{\cos x - \sin x}{\cos x + \sin x}\right), \ 0 < x < \pi$$

Express the matrix as the sum of symmetric and skew symmetric matrix when matrix.

$$\mathbf{A} = \begin{bmatrix} 6 & -2 & 2 \\ -2 & 3 & -1 \\ 2 & -1 & 3 \end{bmatrix}$$

Or

Using the properties of determinants Show that

$$\begin{vmatrix} a-b-c & 2a & 2a \\ 2b & b-c-a & 2b \\ 2c & 2c & c-a-b \end{vmatrix} = (a+b+c)^3$$

17. Differentiate  $x^{sinx}$ , x > 0 w.r.t.x

4

Or

If 
$$e^{y}(x+1) = 1$$
 Show that  $\frac{d^{2}y}{dx^{2}} = \left(\frac{dy}{dx}\right)$ 

18. Evaluate:  $\int \frac{6x+7}{(x-5)(x-4)} dx$ 

4

Or

Evaluate: 
$$\int x^2 \log x \, dx$$

9

- 19. By using properties of definite integrals.
  20. Solve the differential equation
  (e<sup>x</sup> + e<sup>-x</sup>) dy (e<sup>x</sup> e<sup>-x</sup>) dx = 0

  Or

  Find the general solution of the differential equation.
  21. Let a = î + 4ĵ + 2k̂, b = 3î 2ĵ + 7 k̂ & c = 2î ĵ + 4k̂. Find the vector d which is perpendicular to both a and b and c. d = 15. 4

  Or

  Find | x | if for a unit vector a
  (x a).(x + a) = 12

  22. Find the probability of getting 5 exactly twice in 7 throws of a die. 4

  Or

  Find the probability distribution of no. of heads in 4-tosses of a coin
  - Find the area of the region bounded by the ellipse  $\frac{x^2}{4} + \frac{y^2}{9} = 1$

23. Using matrix method, solve following system of linear equations. 6

24. Find the area of the region bounded by the two parabolas  $z=x^2$  and

3x-2y+3z=0; 2x+y-z=1; 4x-3y+2z=4

 $Z^2 = X$ 

25. Find the shortest distance between lines.  $l_1$  and  $l_2$  given by

10

$$\begin{split} \vec{r} &= \hat{i} + 2\hat{j} - 4\hat{k} + \lambda \Big(2\hat{i} + 3\hat{j} + 6\hat{k}\Big) \\ \vec{r} &= 3\hat{i} + 3\hat{j} - 5\hat{k} + \mu \Big(2\hat{i} + 3\hat{j} + 6\hat{k}\Big) \end{split}$$

Or

Find the coordinates of the point where the line through (3, -4, -5) and (2, -3, 1) crosses the plane 2x + y + z = 7

26. Show that of all the rectangles inscribed in a given fixed circle, the square has the maximum area. 6

Or

Find the equations of the narmals to the curve  $z=x^3+2x+6$  which are parallel to the line x+14y+4=0

27. Minimize z = 200x + 500ysubject to the constraints 6

$$x + 2y \ge 10$$
  
 $3x + 4y \le 24$   
 $x \ge 0, y \ge 0$