



4.  $\int a^x dx$  equals to

(a)  $a^x - \log a + c$  (b)  $\frac{\log a}{a^x} + c$

(c)  $\frac{a^x}{\log a} + c$  (d)  $a^x + c$

5.  $\int x^2 e^{x^3} dx$  equals to

(a)  $\frac{1}{3} x e^{x^3}$  (b)  $\frac{1}{3} e^{x^3} + c$  (c)  $\frac{1}{2} e^{x^3} + c$  (d) None of these

6.  $\int x \cdot \sin x dx$  equals.

(a)  $-x \sin x + \cos x + c$  (b)  $-x \cos x + \sin x + c$

(c)  $-x \cos x - \sin x + c$  (d)  $\sin x + \cos x + c$

7.  $\int \frac{(\log x)^2}{x} dx$  equals

(a)  $\frac{1}{3} (\log |x|)^3 + c$  (b)  $\frac{\log x}{x}$

(c)  $\frac{(\log x)^3}{x^3}$  (d) None of these

8.  $\int \frac{dx}{\sin^2 x \cos^2 x}$  equals

- (a)  $\tan x + \cot x + c$                       (b)  $\tan x - \cot x + c$   
 (c)  $\tan x \cot x + c$                       (d)  $\tan x - \cot 2x + c$
9.  $\int \sec x \, dx$  equals to  
 (a)  $\log |\sec x + \tan x| + c$                       (b)  $\sec x - \tan x + c$   
 (c)  $\log |\sec x - \tan x| + c$                       (d)  $\log |\operatorname{cosec} x - \cot x| + c$
10. Area of the region bounded by the curve  $y^2 = 4x$ ,  $y$ -axis and the line  $y = 3$  is  
 (a) 2                      (b)  $\frac{9}{4}$   
 (c)  $\frac{9}{5}$                       (d)  $\frac{9}{2}$
11. The order of the differential equation  $2x^2 \frac{d^2y}{dx^2} - \frac{3dy}{dx} + y = 0$  is  
 (a) 2                      (b) 1  
 (c) 0                      (d) Not defined
12. The general solution of the differential equation  $\frac{dy}{dx} = e^{x+y}$  is  
 (a)  $e^x + e^{-y} = c$                       (b)  $e^x + e^y = c$   
 (c)  $e^{-x} + e^y = c$                       (d)  $e^{-x} + e^{-y} = c$

13. The number of arbitrary constants in the General solution of a differential equation of fourth order are

- (a) 0 (b) 2 (c) 3 (d) 4

14. The angle between two non-zero vector  $\vec{a}$  and  $\vec{b}$  is given by

(a)  $\sin \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$  (b)  $\sin \theta = \frac{|\vec{a} \times \vec{b}|}{|\vec{a}| |\vec{b}|}$

(c)  $\sin \theta = \frac{|\vec{a} \cdot \vec{b}|}{|\vec{a}| |\vec{b}|}$  (d) None of these

15. What is the projection of the vector  $\hat{i} - \hat{j}$  the vector  $\hat{i} + \hat{j}$

- (a) 0 (b) 1 (c) 2 (d) 3

16. The value of  $|\vec{a} \times \vec{b}|$  if  $\vec{a} = 2\hat{i} + 2\hat{j} + 3\hat{k}$  and  $\vec{b} = 3\hat{i} + 5\hat{j} - 2\hat{k}$

- (a)  $\sqrt{705}$  (b)  $\sqrt{507}$   
(c) 507 (d) 0

17. If a line has direction ratio 2, -1, -2 Then the direction cosines are

- (a)  $\frac{3}{2}, \frac{-1}{3}, \frac{-2}{3}$  (b)  $-\frac{1}{3}, \frac{2}{3}, \frac{-2}{3}$   
(c)  $\frac{2}{3}, \frac{-1}{3}, \frac{-2}{3}$  (d) None of these

18. What is the equation of the plane with intercepts 2, 3 and 4 on the x, y and z – axis respectively.

(a)  $\frac{x}{2} + \frac{y}{3} + \frac{z}{4} = 1$       (b)  $\frac{x}{2} + \frac{y}{3} + \frac{z}{4} = 0$

(c)  $2x + 3y + 4z = 1$       (d)  $\frac{x}{4} + \frac{y}{3} + \frac{z}{2} = 1$

19. If  $P(A) = \frac{1}{2}$ ,  $P(B) = 0$ , Then  $P(A/B)$  is

(a)  $= 0$       (b)  $\frac{1}{2}$

(c) Not defined      (d) 1

20. The probability of obtaining an even prime number on each dice, when a pair of dice is called as

(a) 0      (b)  $\frac{1}{3}$

(c)  $\frac{1}{12}$       (d)  $\frac{1}{36}$

21. Find the equation of the tangent to the curve at the pts.

$y = x^4 - 6x^3 + 13x^2 - 10x + 5$  at  $(0, 5)$

OR

Find the equation of the normal at the point  $(am^2, am^3)$  for the curve  $ay^2 = x^3$

22. Integrate the functions

$$(a) \int \frac{x^3 \sin(\tan^{-1} x^4)}{1+x^8} dx \quad \text{OR} \quad \int \frac{3x-1}{(x-1)(x-2)(x-3)} dx$$

$$(b) \int_0^{\pi/2} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx \quad \text{OR} \quad \int e^x (\sin x + \cos x) dx$$

23. Find area of the region bounded by ellipse  $\frac{x^2}{16} + \frac{y^2}{9} = 1$

OR

Find the area bounded by the curve  $(x-1)^2 + y^2 = 1$  and  $x^2 + y^2 = 1$

24. Find the general solution of the differential equation

$$\frac{dy}{dx} = (1+x^2)(1+y^2)$$

OR

$$(x^2 + xy) dy = (x^2 + y^2) dx$$

25. If  $\vec{a}, \vec{b}, \vec{c}$  are unit vector such that  $\vec{a} + \vec{b} + \vec{c} = 0$

find the value of  $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$  is

OR

Find the area of the parallelogram whose adjacent sides are deter-

mined by the vectors  $\vec{a} = \hat{i} - \hat{j}$  and  $\vec{b} = 2\hat{i} - 7\hat{j} + \hat{k}$

26. A die is thrown 6 times. If “getting an odd number” is a success, what is the probability of

(a) 5 success (b) at most 5 success

OR

If  $P(A) = \frac{3}{5}$  and  $P(B) = \frac{1}{5}$ , find  $P(A \cap B)$ , If A and B are independent events.

27. Find the shortest distance between the lines

$$\vec{r} = (\hat{i} + 2\hat{j} + \hat{k}) + \lambda(\hat{i} + \hat{j} + \hat{u}) \text{ and}$$

$$\vec{r} = (2\hat{i} - \hat{j} - \hat{k}) + \mu(2\hat{i} + \hat{j} + 2\hat{k})$$

28. Solve the linear programming problem graphically Maximize

$$Z = 5x + 3y$$

Subject to  $3x + 5y \leq 15$ ,  $5x + 2y \leq 10$   $x \geq 0$   $y \geq 0$