Model Question Paper

Class-XII (Session: 2020-21)(SOS)

Subject-Mathematics

Time Allowed: 3 hrs Maximum Marks: 100 **Special Instructions:-**Same as that of Previous Years Annual question paper (SOS) March 2020. (i) Question numbers 1 to 10 are multiple choice questions of 1 mark each. Q. no. 11 to 14 are 3 marks each. Q. no. 15 to 26 are of 4 marks each and Q. no. 27 to 31 and of 6 marks each. (ii) All questions are compulsory. (iii) 30% more internal choices have been provided from 70% of the syllabus, as 30% syllabus has been deleted due to COVID-19 Pandemic for the session 2020-21 only. 1. If $Sin^{-1} x = y$ then 1 (b) $\frac{-\pi}{2} \le y \le \frac{\pi}{2}$ (a) $0 \le y \le \pi$ (d) $0 < y < \pi$ (c) $0 \le y \le \pi$ 2. If A is symmetric as well as skew symmetric matrix then 1 (a) A is diagonal matrix (b) A is zero matrix (c) A is unit matrix (d) none of these. 3. The derivative of Sec⁻¹ x is (b) $\frac{-1}{|x|\sqrt{1-x^2}}$ (a) $\frac{1}{|x|\sqrt{1-x^2}}$ $(c) \quad \frac{1}{|x|\sqrt{x^2-1}}$

(d) none of these

4. The slope of normal to the curve $y=2x^2+2\sin x$ at x=0 is 1 (a) 3 (b) $\frac{1}{3}$ (c) -3 (d) $\frac{-1}{3}$ 5. $\int \frac{dx}{\sqrt{9x-4x^2}}$ equal to

(a)
$$\frac{1}{9}\sin^{-1}\left(\frac{9x-8}{8}\right) + c$$
 (b) $\frac{1}{2}\sin^{-1}\left(\frac{8x-9}{9}\right) + c$ (c) $\frac{1}{3}\sin^{-1}\left(\frac{9x-8}{8}\right) + c$ (d) $\frac{1}{2}\sin^{-1}\left(\frac{9x-8}{8}\right) + c$

6. The number of arbitrary constants in particular solution of a differential

- equation of fourth order are
 (a) 3 (b) 4 (c) 0 (d) 2

 7. The value of $\hat{i}(\hat{j} \times \hat{k}) + \hat{j}(\hat{i} \times \hat{k}) + \hat{k}(\hat{i} \times \hat{j})$ (a) 0 (b) -1 (c) 1 (d) 3
- 8. If \vec{a} is a non zero vector of magnitude a and λ , a non zero scalar, then $\lambda \vec{a}$ is a unit vector if

(a)
$$\lambda = 1$$
 (b) $\lambda = -1$ (c) $a = |\lambda|$ (d) $a = \frac{1}{|\lambda|}$

9. If a line has direction ratios (2,-1,-2) then its direction cosines will be

(a)
$$<\frac{2}{3}, \frac{-1}{3}, \frac{-2}{3}>$$
 (b) $<\frac{-1}{3}, \frac{2}{3}, \frac{-2}{3}>$

(c)
$$<\frac{-2}{3},\frac{2}{3},\frac{1}{3}>$$
 (d) NOT

- 10. If $P(A) = \frac{6}{11}$, $P(B) = \frac{5}{11}$ and $P(A \cup B) = \frac{7}{11}$ then $P(A \cap B)$ is 1

 - (a) $\frac{6}{11}$ (b) $\frac{7}{11}$ (c) $\frac{5}{11}$ (d) $\frac{4}{11}$

4

11. Find x an y if $2\begin{bmatrix} 1 & 3 \\ 0 & x \end{bmatrix} + \begin{bmatrix} y & 0 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 5 & 6 \\ 1 & 8 \end{bmatrix}$ 3

If A' =
$$\begin{bmatrix} -2 & 3 \\ 1 & 2 \end{bmatrix}$$
, B = $\begin{bmatrix} -1 & 0 \\ 1 & 2 \end{bmatrix}$ then find (A+2B)'

12. Find the value of K so that the function f is continuous at indicated point. 3

$$f(x) = \begin{cases} kx + 1 & x \le 5 \\ 3x - 5 & x > 5 \end{cases} \text{ at } x = 5$$

Differentiate $Cos(\sqrt{x})$ w.r.t x.

- 13. Find the interval in which the given function is strictly increasing or decreasing for $f(x) = 6 - 9x - x^2$
- 14. Find the general solution of the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} + \frac{y}{x} = x^2$$

Or

Find the general solution of differential equation $\sec^2 x \tan y dx + \sec^2 y \tan x dy = 0$

15. Show that the relation R in the set $\{1,2,3\}$ given by $R = \{(1,2),(2,1)\}$ is symmetric but neither reflexive nor transitive.

Find g of & fog if.

$$f(x) = 8x^3$$
 and $g(x) = \frac{1}{x^3}$

16. Prove that $\cos^{-1}\frac{12}{13} + \sin^{-1}\frac{3}{5} = \sin^{-1}\frac{56}{65}$

Express $\tan^{-1} \left(\frac{\cos x}{1 - \sin x} \right) - \frac{3\pi}{2}$ in the simplest form.

17. By using the properties of determinants prove that

$$\begin{vmatrix} 1 & x & x^2 \\ x^2 & 1 & x \\ x & x^2 & 1 \end{vmatrix} = (1 - x^3)^2$$

Or

4

If
$$A = \begin{bmatrix} 3 & 7 \\ 2 & 5 \end{bmatrix}$$
 and $B = \begin{bmatrix} 6 & 8 \\ 7 & 9 \end{bmatrix}$ verify $(AB)^{-1} = B^{-1} A^{-1}$

18. Find $\frac{dy}{dx}$ of $(\log x)^{\cos x}$

Or

Find
$$\frac{dy}{dx}$$
, if $x = at^2$ and $y = 2at$.

19. Evaluate $\int \frac{1}{1+\tan x} dx$

4

4

Evaluate:-
$$\int \frac{1}{\sqrt{x^2 + 2x + 2}} \, dx$$

20. Evaluate: $\int \frac{x}{(x-1)(x-2)(x-3)} dx$

Evaluate:
$$\int \frac{xe^x}{(1+x)^2} dx$$

21. Using properties of definite intergral.

Evaluate:
$$\int_{0}^{\frac{\pi}{2}} \frac{\sin^{4} \times dx}{\sin^{4} x + \cos^{4} x}$$

22. Find the general solution of the differential. equation $(x^2 - y^2) dx + 2xy dy = 0$

Or

Find the particular solution of the differential equation

$$\frac{dy}{dx} + 2y \tan x = \sin x$$
, $y = 0$ where $x = \frac{\pi}{3}$

23. If $\vec{a} = 2\hat{i} + 2\hat{j} + 3\hat{k}$, $\vec{b} = -\hat{i} + 2\hat{j} + \hat{k}$ and $\vec{c} = 3\hat{i} + \hat{j}$ are such that $\vec{a} + \lambda \vec{b}$ is perpendicular to \vec{c} then find the value of λ .

Or

Find the area of parallelogram whose adjacent sides are given by vectors $\vec{a}=3\hat{i}+\hat{j}+4\hat{k}$ and $\vec{b}=\hat{i}-\hat{j}+\hat{k}$

6

$$\frac{1-x}{3} = \frac{7y-14}{2p} = \frac{z-3}{2}$$
 and
$$\frac{7-7x}{3p} = \frac{y-5}{1} = \frac{6-z}{5}$$
 are at right angles.

Or

Find the equation of the plane with intercepts 2,3 and 4 on the x,y,z-axis respectively.

- 25. A die is thrown. If E is the event 'the number appearing is multiple of 3 and 4' F be the event the number appearing is even then find whether E and F are independent?
 4
- 26. From a lot of 30 bulbs which include 6-defective, a sample of 4-bulbs in drawn at random with replacement. Find the probability distribution of the number of defective bulbs.

Or

Find the probability of throwing at most 2 sixes in 6 throws of a single die.

- 27. Solve the following system of linear equation using matrix method. 6 2x 3y + 5z = 11, 3x + 2y 4z = -5; x + y 2z = -3.
- 28. Find two positive number x & y such that their sum is 35 and the product x²y² is maximum.

Or

Prove that the curve $5x = y^2$ and xy=k cut at right angle it $8k^2=1$.

29. Using the intergration find the area of region bounded by the triangle whose vertices are (1,0), (2,2) and (3,1).

Oı

Find the smaller area enclosed by the circle. $x^2 + y^2 = 4$ and the line x + y = 2.

30. Find the shortest distance between the lines

$$\frac{x+1}{7} = \frac{y+1}{-6} = \frac{z+1}{1} \text{ and } \frac{x-3}{1} = \frac{y-5}{-2} = \frac{z-7}{1}$$
Or

Find the equation of plane passing through the point (-1,3,2) and perpendicular to each of the planes x + 2y + 3z = 5 and 3x + 3y + z = 0

31. Maximize z = 3x + 2y

Subject to constraints

$$x + 2y \le 10$$

$$3x + y \le 15$$